

## $\Upsilon$ production in dAu collisions at RHIC

**N. Matagne**  
University of Mons

**Quarkonium Working Group Workshop**  
**May 19, 2010**

In collaboration with E. G. Ferreiro, F. Fleuret, J. P. Lansberg and A.  
Rakotozafindrabe

# Outline

- 1 Introduction and motivations
- 2 Experimental situation
- 3 On the kinematics of  $\Upsilon$  production
- 4 The Glauber Monte Carlo
- 5 Results for dAu collisions for  $\Upsilon$
- 6 EMC effect for gluons
- 7 Conclusions and perspectives

# Introduction and motivations

- Extend to  $\Upsilon$  the study of **CNM effects** (shadowing + absorption) on **production** of quarkonia

# Introduction and motivations

- Extend to  $\Upsilon$  the study of CNM effects (shadowing + absorption) on production of quarkonia
- Glauber Monte Carlo model to simulate dAu and AuAu collisions at RHIC

# Introduction and motivations

- Extend to  $\Upsilon$  the study of **CNM effects** (shadowing + absorption) on **production** of quarkonia
- Glauber Monte Carlo model to simulate **dAu and AuAu collisions at RHIC**
- **Two main production schemes** ( $2 \rightarrow 1$ ,  $2 \rightarrow 2$ )

# Introduction and motivations

- Extend to  $\Upsilon$  the study of **CNM effects** (shadowing + absorption) on **production** of quarkonia
- Glauber Monte Carlo model to simulate **dAu and AuAu collisions at RHIC**
- **Two main production schemes** ( $2 \rightarrow 1$ ,  $2 \rightarrow 2$ )
- Comparison of **three different** shadowing parametrisations


# Introduction and motivations

- Extend to  $\Upsilon$  the study of **CNM effects** (shadowing + absorption) on **production** of quarkonia
- Glauber Monte Carlo model to simulate **dAu and AuAu collisions at RHIC**
- **Two main production schemes** ( $2 \rightarrow 1$ ,  $2 \rightarrow 2$ )
- Comparison of **three different** shadowing parametrisations
- **Three** absorption cross sections

# Experimental situation

P. Artoisenet, J. Campbell, J.P. Lansberg, F. Maltoni, Phys. Rev. Lett. 101, 152001 (2008).  
D. Acosta et al. (CDF collaboration), Phys. Rev. Lett 88, 161802 (2002).

## Results at 1.8 TeV




- CSM describes well the data at NNLO\*



# Experimental situation

P. Artoisenet, J. Campbell, J.P. Lansberg, F. Maltoni, Phys. Rev. Lett. 101, 152001 (2008).  
D. Acosta et al. (CDF collaboration), Phys. Rev. Lett 88, 161802 (2002).

## Results at 1.8 TeV




- CSM describes well the data at NNLO\*
- However LO CSM is sufficient to describe low  $p_T$  data

# Experimental situation

S. J. Brodsky and J. P. Lansberg, Phys. Rev. D81, 014004 (2010).  
 P. Djawotho et al., J. Phys. G34, s947 (2007); T. Ullrich (private communication) (STAR)  
 C.L. da Silva, Nucl. Phys. A830, 227c (2009); L.L. Levy, Nucl. Phys. A830, 353c (2009); W. Xie et al., J. Phys. A774, 693 (2006) (PHENIX)

## Results at 200 GeV



- Upper dashed line,  $m_b = 4.5$  GeV,  $\mu_R = M_T$ ,  $\mu_F = 2M_T$
- Lower dashed line,  $m_b = 5.0$  GeV,  $\mu_R = 2M_T$ ,  $\mu_F = M_T$

We take the parameters of the upper curve in the following.


# Experimental situation

H. Liua and the STAR collaboration, Nucl. Phys. A830, 235c (2009)

H. Pereira Da Costa for the PHENIX collaboration, talk at the rencontres de Moriond, March 15, 2010

3 data points :

- $-2.2 < y < -1.2$ : PHENIX -  $R_{dAu} = 0.84 \pm 0.34$  (stat.)  $\pm 0.28$  (sys.)
- $1.2 < y < 2.2$ : PHENIX -  $R_{dAu} = 0.53 \pm 0.20$  (stat.)  $\pm 0.16$  (sys.)



- $|y| < 0.5$ : STAR -  $R_{dAu} = 0.98 \pm 0.32$  (stat.)  $\pm 0.28$  (sys.)

# On the kinematics of $\Upsilon$ production

If  $\mathcal{F}_g^A(x, \vec{r}, z, \mu_f)$  gives the **distribution of a gluon** of mom. fract.  $x$  at a **position  $\vec{r}, z$  in a nucleus  $A$** , the differential cross-section reads:

$$\frac{d\sigma_{AB}}{dy dP_T d\vec{b}} =$$

$2 \rightarrow 1$  kinematics with **intrinsic**  $p_T$

$2 \rightarrow 2$  kinematics with **extrinsic**  $p_T$

# On the kinematics of $\Upsilon$ production

If  $\mathcal{F}_g^A(x, \vec{r}, z, \mu_f)$  gives the **distribution of a gluon** of mom. fract.  $x$  at a **position  $\vec{r}, z$  in a nucleus  $A$** , the differential cross-section reads:

$$\frac{d\sigma_{AB}}{dy dP_T d\vec{b}} =$$

**$2 \rightarrow 1$  kinematics with intrinsic  $p_T$**

$$\begin{aligned} & \int d\vec{r}_A dz_A dz_B \\ & \times \mathcal{F}_g^A(x_1^0, \vec{r}_A, z_A, \mu_f) \mathcal{F}_g^B(x_2^0, \vec{r}_B, z_B, \mu_f) \\ & \times \sigma_{gg}^{\text{Intr.}}(x_1^0, x_2^0) \\ & \times S_A(\vec{r}_A, z_A) S_B(\vec{r}_B, z_B) \end{aligned}$$

**$2 \rightarrow 2$  kinematics with extrinsic  $p_T$**

$$\begin{aligned} & \int dx_1 dx_2 \int d\vec{r}_A dz_A dz_B \\ & \times \mathcal{F}_g^A(x_1, \vec{r}_A, z_A, \mu_f) \mathcal{F}_g^B(x_2, \vec{r}_B, z_B, \mu_f) \\ & \times 2\hat{s} P_T \frac{d\sigma_{gg \rightarrow \Upsilon + g}}{d\hat{t}} \delta(\hat{s} - \hat{t} - \hat{u} - M^2) \\ & \times S_A(\vec{r}, z_A) S_B(\vec{r}_B, z_B) \end{aligned}$$

# On the kinematics of $\Upsilon$ production

If  $\mathcal{F}_g^A(x, \vec{r}, z, \mu_f)$  gives the **distribution of a gluon** of mom. fract.  $x$  at a **position  $\vec{r}, z$  in a nucleus  $A$** , the differential cross-section reads:

$$\frac{d\sigma_{AB}}{dy dP_T d\vec{b}} =$$

**$2 \rightarrow 1$  kinematics with intrinsic  $p_T$**

$$\begin{aligned} & \int d\vec{r}_A dz_A dz_B \\ & \times \mathcal{F}_g^A(x_1^0, \vec{r}_A, z_A, \mu_f) \mathcal{F}_g^B(x_2^0, \vec{r}_B, z_B, \mu_f) \\ & \times \sigma_{gg}^{\text{Intr.}}(x_1^0, x_2^0) \\ & \times S_A(\vec{r}_A, z_A) S_B(\vec{r}_B, z_B) \end{aligned}$$



$$x_{1,2} = \frac{m_T}{\sqrt{s_{NN}}} \exp(\pm y) \equiv x_{1,2}^0(y, P_T)$$

**$2 \rightarrow 2$  kinematics with extrinsic  $p_T$**

$$\begin{aligned} & \int dx_1 dx_2 \int d\vec{r}_A dz_A dz_B \\ & \times \mathcal{F}_g^A(x_1, \vec{r}_A, z_A, \mu_f) \mathcal{F}_g^B(x_2, \vec{r}_B, z_B, \mu_f) \\ & \times 2\hat{s} P_T \frac{d\sigma_{gg \rightarrow \Upsilon+g}}{d\hat{t}} \delta(\hat{s} - \hat{t} - \hat{u} - M^2) \\ & \times S_A(\vec{r}, z_A) S_B(\vec{r}_B, z_B) \end{aligned}$$



$$\delta(\cdot) \rightarrow x_2 = \frac{x_1 m_T \sqrt{s_{NN}} e^{-y} - M^2}{\sqrt{s_{NN}} (\sqrt{s_{NN}} x_1 - m_T e^y)}$$

# On the kinematics of $\Upsilon$ production




For a given couple  $(y, p_T)$ ,  $x_2$  is larger in the extrinsic scheme

# On the kinematics of $\Upsilon$ production





For a given couple  $(y, p_T)$ ,  $x_2$  is larger in the extrinsic scheme

Antishadowing  
peak at  $\sim 10^{-1}$






# On the kinematics of $\Upsilon$ production



Antishadowing  
peak at  $\sim 10^{-1}$


We expect **different shadowing effects**  
in both cases.


For a given couple  
( $y, p_T$ ),  $x_2$  is **larger** in  
the **extrinsic** scheme



# The Glauber Monte Carlo (for $\Upsilon$ here)

## Our Monte-Carlo approach for $J/\Psi$ production



Results for dAu collisions for  $\Upsilon$ 


EKS shadowing :

—■— Intrinsic pT

- - -●- - Extrinsic pT

# Results for dAu collisions for $\Upsilon$

R. Vogt, talk at Joint CATHIE-TECHQM Meeting, BNL, December 14-18, 2009




EKS98 (blue), nDSG (magenta), EPS08 (red), EPS09 (cyan)

- One has to be careful about binning effect (usually decrease the modifications)
- Interesting to see the difference between  $2- > 1$  (as done by R. Vogt) and  $2- > 2$  (kinematics for LO CSM)

# Results for dAu collisions for $\Upsilon$

H. Liua and the STAR collaboration, Nucl. Phys. A830, 235c (2009)


H. Pereira Da Costa for the PHENIX collaboration, talk at the rencontres de Moriond, March 15, 2010



- backward: ok within uncertainties;
- central: reasonable job -  $R_{dAu} > 1$  (for any  $\sigma_{abs}$ );
- forward: clearly too high (for any  $\sigma_{abs}$ );

# Results for dAu collisions for $\Upsilon$

H. Liua and the STAR collaboration, Nucl. Phys. A830, 235c (2009)  
H. Pereira Da Costa for the PHENIX collaboration, talk at the rencontres de Moriond, March 15, 2010




- backward: slightly too high (but ok within uncertainties);
- central: reasonable job -  $R_{dAu} > 1$  (for any  $\sigma_{abs}$ );
- forward: clearly too high (for any  $\sigma_{abs}$ ), though 'better' than EKS;


# Results for dAu collisions for $\Upsilon$

H. Liua and the STAR collaboration, Nucl. Phys. A830, 235c (2009)

H. Pereira Da Costa for the PHENIX collaboration, talk at the rencontres de Moriond, March 15, 2010



- backward: ok within uncertainties;
- central: reasonable job -  $R_{dAu}$  1;
- forward : clearly too high (for any  $\sigma_{abs}$ ), though 'better' than EKS and EPS;


Results for dAu collisions for  $\Upsilon$ 

EKS

The extrinsic scheme enables to predict the  $p_T$  dependence, which is non trivial

- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb




Results for dAu collisions for  $\Upsilon$ 

EPS

The extrinsic scheme enables to predict the  $p_T$  dependence, which is non trivial


- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb

Results for dAu collisions for  $\Upsilon$ 

nDSg


The extrinsic scheme enables to predict the  $p_T$  dependence, which is non trivial

- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb


Results for dAu collisions for  $\Upsilon$ 

EKS

- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb

Results for dAu collisions for  $\Upsilon$ 

- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb


Results for dAu collisions for  $\Upsilon$ 

nDSg

- In blue,  $\sigma_{abs} = 0.0$  mb
- In green,  $\sigma_{abs} = 0.5$  mb
- In red,  $\sigma_{abs} = 1.0$  mb


# EMC effect for gluons

- Tension between the theory and the PHENIX point in the backward region
- The backward region correspond to the EMC region ( $x > 0.1$ )
- EMC effect basically unknown for the gluon



# EMC effect for gluons


- Let us try to increase the suppression of  $g(x)$  in the EMC region
- Keeping momentum conservation :  $\int xg(x) dx = Cst$



# EMC effect for gluons

H. Liua and the STAR collaboration, Nucl. Phys. A830, 235c (2009)

H. Pereira Da Costa for the PHENIX collaboration, talk at the rencontres de Moriond, March 15, 2010



**Works better** for backward region



# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy

# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy
- Only very forward (backward) region shows  $R_{dAU} < 1$  due to shadowing

# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy
- Only very forward (backward) region shows  $R_{dAU} < 1$  due to shadowing
- Backward region dominated by EMC effect

# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy
- Only very forward (backward) region shows  $R_{dAU} < 1$  due to shadowing
- Backward region dominated by EMC effect
- Energy loss seems to be needed to reproduce the data at forward  $y$

# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy
- Only very forward (backward) region shows  $R_{dAU} < 1$  due to shadowing
- Backward region dominated by EMC effect
- Energy loss seems to be needed to reproduce the data at forward  $y$
- Within the commonly accepted  $\sigma_{abs}$ , one should expect an excess of  $\Upsilon$

# Conclusions and perspectives

- Production of Upsilon sits in the antishadowing region at RHIC energy
- Only very forward (backward) region shows  $R_{dAU} < 1$  due to shadowing
- Backward region dominated by EMC effect
- Energy loss seems to be needed to reproduce the data at forward  $y$
- Within the commonly accepted  $\sigma_{abs}$ , one should expect an excess of  $\Upsilon$
- ... unless there is no antishadowing (see nDSg)